



---

# **AP<sup>®</sup> Physics 2: Algebra-Based 2016 Free-Response Questions**

---

© 2016 The College Board. College Board, Advanced Placement Program, AP, AP Central, and the acorn logo are registered trademarks of the College Board.

Visit the College Board on the Web: [www.collegeboard.org](http://www.collegeboard.org).

AP Central is the official online home for the AP Program: [apcentral.collegeboard.org](http://apcentral.collegeboard.org).

## AP<sup>®</sup> PHYSICS 2 TABLE OF INFORMATION

CONSTANTS AND CONVERSION FACTORS	
Proton mass, $m_p = 1.67 \times 10^{-27}$ kg Neutron mass, $m_n = 1.67 \times 10^{-27}$ kg Electron mass, $m_e = 9.11 \times 10^{-31}$ kg Avogadro's number, $N_0 = 6.02 \times 10^{23}$ mol <sup>-1</sup> Universal gas constant, $R = 8.31$ J/(mol·K) Boltzmann's constant, $k_B = 1.38 \times 10^{-23}$ J/K	Electron charge magnitude, $e = 1.60 \times 10^{-19}$ C 1 electron volt, $1 \text{ eV} = 1.60 \times 10^{-19}$ J Speed of light, $c = 3.00 \times 10^8$ m/s Universal gravitational constant, $G = 6.67 \times 10^{-11}$ m <sup>3</sup> /kg·s <sup>2</sup> Acceleration due to gravity at Earth's surface, $g = 9.8$ m/s <sup>2</sup>
1 unified atomic mass unit, Planck's constant, Vacuum permittivity, Coulomb's law constant, $k = 1/4\pi\epsilon_0 = 9.0 \times 10^9$ N·m <sup>2</sup> /C <sup>2</sup> Vacuum permeability, Magnetic constant, $k' = \mu_0/4\pi = 1 \times 10^{-7}$ (T·m)/A 1 atmosphere pressure,	$1 \text{ u} = 1.66 \times 10^{-27}$ kg = 931 MeV/c <sup>2</sup> $h = 6.63 \times 10^{-34}$ J·s = $4.14 \times 10^{-15}$ eV·s $hc = 1.99 \times 10^{-25}$ J·m = $1.24 \times 10^3$ eV·nm $\epsilon_0 = 8.85 \times 10^{-12}$ C <sup>2</sup> /N·m <sup>2</sup> $\mu_0 = 4\pi \times 10^{-7}$ (T·m)/A $1 \text{ atm} = 1.0 \times 10^5$ N/m <sup>2</sup> = $1.0 \times 10^5$ Pa

UNIT SYMBOLS	meter,	m	mole,	mol	watt,	W	farad,	F
	kilogram,	kg	hertz,	Hz	coulomb,	C	tesla,	T
	second,	s	newton,	N	volt,	V	degree Celsius,	°C
	ampere,	A	pascal,	Pa	ohm,	Ω	electron volt,	eV
	kelvin,	K	joule,	J	henry,	H		

PREFIXES		
Factor	Prefix	Symbol
10 <sup>12</sup>	tera	T
10 <sup>9</sup>	giga	G
10 <sup>6</sup>	mega	M
10 <sup>3</sup>	kilo	k
10 <sup>-2</sup>	centi	c
10 <sup>-3</sup>	milli	m
10 <sup>-6</sup>	micro	μ
10 <sup>-9</sup>	nano	n
10 <sup>-12</sup>	pico	p

VALUES OF TRIGONOMETRIC FUNCTIONS FOR COMMON ANGLES							
$\theta$	0°	30°	37°	45°	53°	60°	90°
$\sin \theta$	0	1/2	3/5	$\sqrt{2}/2$	4/5	$\sqrt{3}/2$	1
$\cos \theta$	1	$\sqrt{3}/2$	4/5	$\sqrt{2}/2$	3/5	1/2	0
$\tan \theta$	0	$\sqrt{3}/3$	3/4	1	4/3	$\sqrt{3}$	∞

The following conventions are used in this exam.

- I. The frame of reference of any problem is assumed to be inertial unless otherwise stated.
- II. In all situations, positive work is defined as work done on a system.
- III. The direction of current is conventional current: the direction in which positive charge would drift.
- IV. Assume all batteries and meters are ideal unless otherwise stated.
- V. Assume edge effects for the electric field of a parallel plate capacitor unless otherwise stated.
- VI. For any isolated electrically charged object, the electric potential is defined as zero at infinite distance from the charged object

## AP<sup>®</sup> PHYSICS 2 EQUATIONS

### MECHANICS

$v_x = v_{x0} + a_x t$	$a = \text{acceleration}$
$x = x_0 + v_{x0} t + \frac{1}{2} a_x t^2$	$A = \text{amplitude}$
$v_x^2 = v_{x0}^2 + 2a_x(x - x_0)$	$d = \text{distance}$
$\vec{a} = \frac{\sum \vec{F}}{m} = \frac{\vec{F}_{net}}{m}$	$E = \text{energy}$
$ \vec{F}_f  \leq \mu  \vec{F}_n $	$F = \text{force}$
$a_c = \frac{v^2}{r}$	$f = \text{frequency}$
$\vec{p} = m\vec{v}$	$I = \text{rotational inertia}$
$\Delta\vec{p} = \vec{F} \Delta t$	$K = \text{kinetic energy}$
$K = \frac{1}{2} m v^2$	$k = \text{spring constant}$
$\Delta E = W = F_{\parallel} d = F d \cos \theta$	$L = \text{angular momentum}$
$P = \frac{\Delta E}{\Delta t}$	$\ell = \text{length}$
$\theta = \theta_0 + \omega_0 t + \frac{1}{2} \alpha t^2$	$m = \text{mass}$
$\omega = \omega_0 + \alpha t$	$P = \text{power}$
$x = A \cos(\omega t) = A \cos(2\pi f t)$	$p = \text{momentum}$
$x_{cm} = \frac{\sum m_i x_i}{\sum m_i}$	$r = \text{radius or separation}$
$\vec{\alpha} = \frac{\sum \vec{\tau}}{I} = \frac{\vec{\tau}_{net}}{I}$	$T = \text{period}$
$\tau = r_{\perp} F = r F \sin \theta$	$t = \text{time}$
$L = I \omega$	$U = \text{potential energy}$
$\Delta L = \tau \Delta t$	$v = \text{speed}$
$K = \frac{1}{2} I \omega^2$	$W = \text{work done on a system}$
$ \vec{F}_s  = k  \vec{x} $	$x = \text{position}$
	$y = \text{height}$
	$\alpha = \text{angular acceleration}$
	$\mu = \text{coefficient of friction}$
	$\theta = \text{angle}$
	$\tau = \text{torque}$
	$\omega = \text{angular speed}$
	$U_s = \frac{1}{2} k x^2$
	$\Delta U_g = mg \Delta y$
	$T = \frac{2\pi}{\omega} = \frac{1}{f}$
	$T_s = 2\pi \sqrt{\frac{m}{k}}$
	$T_p = 2\pi \sqrt{\frac{\ell}{g}}$
	$ \vec{F}_g  = G \frac{m_1 m_2}{r^2}$
	$\vec{g} = \frac{\vec{F}_g}{m}$
	$U_G = -\frac{G m_1 m_2}{r}$

### ELECTRICITY AND MAGNETISM

$ \vec{F}_E  = \frac{1}{4\pi\epsilon_0} \frac{ q_1 q_2 }{r^2}$	$A = \text{area}$
$\vec{E} = \frac{\vec{F}_E}{q}$	$B = \text{magnetic field}$
$ \vec{E}  = \frac{1}{4\pi\epsilon_0} \frac{ q }{r^2}$	$C = \text{capacitance}$
$\Delta U_E = q \Delta V$	$d = \text{distance}$
$V = \frac{1}{4\pi\epsilon_0} \frac{q}{r}$	$E = \text{electric field}$
$ \vec{E}  = \left  \frac{\Delta V}{\Delta r} \right $	$\mathcal{E} = \text{emf}$
$\Delta V = \frac{Q}{C}$	$F = \text{force}$
$C = \kappa \epsilon_0 \frac{A}{d}$	$I = \text{current}$
$E = \frac{Q}{\epsilon_0 A}$	$\ell = \text{length}$
$U_C = \frac{1}{2} Q \Delta V = \frac{1}{2} C (\Delta V)^2$	$P = \text{power}$
$I = \frac{\Delta Q}{\Delta t}$	$Q = \text{charge}$
$R = \frac{\rho \ell}{A}$	$q = \text{point charge}$
$P = I \Delta V$	$R = \text{resistance}$
$I = \frac{\Delta V}{R}$	$r = \text{separation}$
$R_s = \sum_i R_i$	$t = \text{time}$
$\frac{1}{R_p} = \sum_i \frac{1}{R_i}$	$U = \text{potential (stored) energy}$
$C_p = \sum_i C_i$	$V = \text{electric potential}$
$\frac{1}{C_s} = \sum_i \frac{1}{C_i}$	$v = \text{speed}$
$B = \frac{\mu_0 I}{2\pi r}$	$\kappa = \text{dielectric constant}$
	$\rho = \text{resistivity}$
	$\theta = \text{angle}$
	$\Phi = \text{flux}$
	$\vec{F}_M = q\vec{v} \times \vec{B}$
	$ \vec{F}_M  =  q\vec{v}   \sin \theta   \vec{B} $
	$\vec{F}_M = I\vec{\ell} \times \vec{B}$
	$ \vec{F}_M  =  I\vec{\ell}   \sin \theta   \vec{B} $
	$\Phi_B = \vec{B} \cdot \vec{A}$
	$\Phi_B =  \vec{B}  \cos \theta  \vec{A} $
	$\mathcal{E} = -\frac{\Delta \Phi_B}{\Delta t}$
	$\mathcal{E} = B l v$

## AP<sup>®</sup> PHYSICS 2 EQUATIONS

### FLUID MECHANICS AND THERMAL PHYSICS

$$\rho = \frac{m}{V}$$

$$P = \frac{F}{A}$$

$$P = P_0 + \rho gh$$

$$F_b = \rho Vg$$

$$A_1 v_1 = A_2 v_2$$

$$P_1 + \rho g y_1 + \frac{1}{2} \rho v_1^2 = P_2 + \rho g y_2 + \frac{1}{2} \rho v_2^2$$

$$\frac{Q}{\Delta t} = \frac{kA \Delta T}{L}$$

$$PV = nRT = Nk_B T$$

$$K = \frac{3}{2} k_B T$$

$$W = -P \Delta V$$

$$\Delta U = Q + W$$

*A* = area  
*F* = force  
*h* = depth  
*k* = thermal conductivity  
*K* = kinetic energy  
*L* = thickness  
*m* = mass  
*n* = number of moles  
*N* = number of molecules  
*P* = pressure  
*Q* = energy transferred to a system by heating  
*T* = temperature  
*t* = time  
*U* = internal energy  
*V* = volume  
*v* = speed  
*W* = work done on a system  
*y* = height  
*ρ* = density

### MODERN PHYSICS

$$E = hf$$

$$K_{\max} = hf - \phi$$

$$\lambda = \frac{h}{p}$$

$$E = mc^2$$

*E* = energy  
*f* = frequency  
*K* = kinetic energy  
*m* = mass  
*p* = momentum  
*λ* = wavelength  
*φ* = work function

### WAVES AND OPTICS

$$\lambda = \frac{v}{f}$$

$$n = \frac{c}{v}$$

$$n_1 \sin \theta_1 = n_2 \sin \theta_2$$

$$\frac{1}{s_i} + \frac{1}{s_o} = \frac{1}{f}$$

$$|M| = \left| \frac{h_i}{h_o} \right| = \left| \frac{s_i}{s_o} \right|$$

$$\Delta L = m\lambda$$

$$d \sin \theta = m\lambda$$

*d* = separation  
*f* = frequency or focal length  
*h* = height  
*L* = distance  
*M* = magnification  
*m* = an integer  
*n* = index of refraction  
*s* = distance  
*v* = speed  
*λ* = wavelength  
*θ* = angle

### GEOMETRY AND TRIGONOMETRY

Rectangle  
 $A = bh$

Triangle  
 $A = \frac{1}{2}bh$

Circle  
 $A = \pi r^2$   
 $C = 2\pi r$

Rectangular solid  
 $V = \ell wh$

Cylinder  
 $V = \pi r^2 \ell$   
 $S = 2\pi r \ell + 2\pi r^2$

Sphere  
 $V = \frac{4}{3}\pi r^3$   
 $S = 4\pi r^2$


*A* = area  
*C* = circumference  
*V* = volume  
*S* = surface area  
*b* = base  
*h* = height  
*ℓ* = length  
*w* = width  
*r* = radius

Right triangle  
 $c^2 = a^2 + b^2$

$$\sin \theta = \frac{a}{c}$$


$$\cos \theta = \frac{b}{c}$$

$$\tan \theta = \frac{a}{b}$$



**2016 AP<sup>®</sup> PHYSICS 2 FREE-RESPONSE QUESTIONS****PHYSICS 2****Section II****4 Questions****Time—90 minutes**

**Directions:** Questions 1 and 4 are short free-response questions that require about 20 minutes each to answer and are worth 10 points each. Questions 2 and 3 are long free-response questions that require about 25 minutes each to answer and are worth 12 points each. Show your work for each part in the space provided after that part.



1. (10 points, suggested time 20 minutes)

Two moles of a monatomic ideal gas are enclosed in a cylinder by a movable piston. The gas is taken through the thermodynamic cycle shown in the figure above. The piston has a cross-sectional area of  $5 \times 10^{-3} \text{ m}^2$ .

(a)

- Calculate the force that the gas exerts on the piston in state *A*, and explain how the collisions of the gas atoms with the piston allow the gas to exert a force on the piston.
- Calculate the temperature of the gas in state *B*, and indicate the microscopic property of the gas that is characterized by the temperature.

(b)

- Predict qualitatively how the internal energy of the gas changes as it is taken from state *A* to state *B*. Justify your prediction.
- Calculate the energy added to the gas by heating as it is taken from state *A* to state *C* along the path *ABC*.

(c) Determine the change in the total kinetic energy of the gas atoms as the gas is taken directly from state *C* to state *A*.


## 2016 AP<sup>®</sup> PHYSICS 2 FREE-RESPONSE QUESTIONS

2. (12 points, suggested time 25 minutes)

A student is given a glass block that has been specially treated so that the path of light can be seen as the light travels through the glass. The student is asked to design an experiment to measure the index of refraction of the glass. The light source available in the laboratory is a hydrogen lamp that emits red light of a known wavelength.

- (a) A linear graph is to be used to determine the index of refraction of the glass. Indicate the quantities that should be graphed and describe how the graph could be used to determine the index of refraction of the glass.
- (b) Outline an experimental procedure that could gather the necessary data. Include sufficient detail so that another student could follow your procedure. In addition to the glass block and the hydrogen lamp, the equipment in a typical classroom laboratory is available.
- (c) Predict how the path of the light will change as it enters the glass. Support your prediction using a qualitative comparison of the speed of light in glass and the speed of light in air.
- (d) Describe the process(es) by which red light from the lamp is produced by hydrogen atoms that are initially in the ground state. Draw and label an energy level diagram that supports the atomic process(es) you describe.

2016 AP<sup>®</sup> PHYSICS 2 FREE-RESPONSE QUESTIONS



3. (12 points, suggested time 25 minutes)

The dots in the figure above represent two identical spheres,  $X$  and  $Y$ , that are fixed in place with their centers in the plane of the page. Both spheres are charged, and the charge on sphere  $Y$  is positive. The lines are isolines of electric potential, also in the plane of the page, with a potential difference of 10 V between each set of adjacent lines. The absolute value of the electric potential of the outermost line is 50 V.

(a) Indicate the values of the potentials, including the signs, at the labeled points  $A$  and  $B$ .

Potential at point  $A$  \_\_\_\_\_ Potential at point  $B$  \_\_\_\_\_

(b)

- i. How do the magnitudes and the signs of the charges of the spheres compare? Explain your answer in terms of the isolines of electric potential shown.
- ii. The spheres at points  $X$  and  $Y$  have masses in the same ratio as the magnitudes of their charges. The isolines of gravitational potential for the spheres have shapes similar to those of the isolines shown. Explain why the two sets of isolines have similar shapes.

Let the potentials at the three labeled points be  $V_A$ ,  $V_B$ , and  $V_C$ . A proton with charge  $+q$  and mass  $m$  is released from rest at point  $B$ .

(c) Based on your answer to part (b)(ii), briefly describe one similarity and one difference between the electric and gravitational forces exerted on the proton by the system of the two spheres. The similarity and difference you describe must not be ones that generally apply to all forces.

## 2016 AP<sup>®</sup> PHYSICS 2 FREE-RESPONSE QUESTIONS

- (d) At some time after being released from rest at point  $B$ , the proton has moved through a potential difference of magnitude 20 V.
- Determine the change in electric potential energy of the proton-spheres system when the proton has moved through the 20 V potential difference. Express your answer symbolically in terms of  $q$ ,  $V_A$ ,  $V_B$ ,  $V_C$ , and physical constants, as appropriate.
  - As it moved through the 20 V potential difference, the proton was displaced a distance  $d$  by the electric force. Determine a symbolic expression for the total work done on the proton by the electric field in terms of the average magnitude  $E_{avg}$  of the electric field over that distance.
  - Two students are discussing how and why the kinetic energy of the proton would change after it is released.
    - Student 1 says that if the system is defined as the proton and the spheres, the increase in the proton's kinetic energy is due to a change in the system's potential energy as the proton moves through the 20 V potential difference.
    - Student 2 says that if the system is defined as only the proton, the kinetic energy of the proton increases because positive work is done on the proton by the electric field as the proton moves through the 20 V potential difference.Discuss each student's claims, explaining why each is correct or incorrect.



## 2016 AP<sup>®</sup> PHYSICS 2 FREE-RESPONSE QUESTIONS

4. (10 points, suggested time 20 minutes)

Some students are investigating the behavior of a circuit with four components in series: a resistor of resistance  $R$ , a capacitor of capacitance  $C$ , a battery with potential difference  $\mathcal{E}$ , and a switch. Initially, the capacitor is uncharged and the switch is open.

(a)

- i. Determine the current in the resistor and the potential difference across the capacitor immediately after the switch is closed.
- ii. Determine the current in the resistor and the potential difference across the capacitor a long time after the switch is closed.

(b) The switch is opened, the capacitor is discharged, and a second, identical capacitor is added to the circuit in series with the other components. The switch is then closed again.

- i. A long time after the switch is closed, the energy stored in the single capacitor in the original circuit is  $U_1$ , and the total energy stored in the two capacitors in the new circuit is  $U_2$ . Calculate the ratio  $U_1/U_2$ .
- ii. The two capacitors in series are to be replaced with a single capacitor that will have the same energy  $U_2$ . Indicate a plate area and a distance between the plates for the new capacitor, compared with one of the original capacitors, that will accomplish this. Support your reasoning using appropriate physics principles and/or mathematical models.

The students are then asked to design two circuits each containing a switch, a battery with a small internal resistance, a lightbulb, and a capacitor. In arrangement 1, the bulb should gradually light up after the switch is closed, becoming brightest after the switch has been closed a long time. In arrangement 2, the bulb should be brightest when the switch is first closed, getting dimmer with time, and going out completely when the switch has been closed for a long time.

(c) Using standard symbols, draw two circuit diagrams, one showing a possible circuit for arrangement 1 and the other showing a possible circuit for arrangement 2. Justify your circuit diagrams with a paragraph-length explanation referring to the properties of lightbulbs and capacitors in circuits and the conservation of energy and/or the conservation of charge.

**STOP**

**END OF EXAM**